# FENTANYL<sup>\*</sup>

## SAFETY RECOMMENDATIONS FOR FIRST RESPONDERS

- <sup>†</sup> For the purposes of this document, fentanyl, related substances, and synthetic opioids (herein after referred to as fentanyl<sup>†</sup>) includes fentanyl analogues (e.g., acetylfentanyl, acrylfentanyl, carfentanil, furanylfentanyl), novel synthetic opioids (e.g., U-47700), and other drugs that may be laced with these substances.
- The abuse of drugs containing fentanyl<sup>†</sup> is killing Americans. Misinformation and inconsistent recommendations regarding fentanyl<sup>†</sup> have resulted in confusion in the first responder community.
- ▶ You as a first responder (law enforcement, fire, rescue, and emergency medical services (EMS) personnel) are increasingly likely to encounter fentanyl<sup>†</sup> in your daily activities (e.g., responding to overdose calls, conducting traffic stops, arrests, and searches).
- This document provides scientific, evidence-based recommendations to protect yourself from exposure.

### WHAT YOU NEED TO KNOW

- Fentanyl<sup>+</sup> can be present in a variety of forms (e.g., powder, tablets, capsules, solutions, and rocks).
- Inhalation of airborne powder is MOST LIKELY to lead to harmful effects, but is less likely to occur than skin contact.
- Incidental skin contact may occur during daily activities but is not expected to lead to harmful effects if the contaminated skin is promptly washed off with water.
- Personal Protective Equipment (PPE) is effective in protecting you from exposure.
- Slow breathing or no breathing, drowsiness or unresponsiveness, and constricted or pinpoint pupils are the specific signs consistent with fentanyl<sup>†</sup> intoxication.
- Naloxone is an effective medication that rapidly reverses the effects of fentanyl<sup>†</sup>.

# To protect yourself from exposure

- Wear gloves when the presence of fentanyl<sup>†</sup> is suspected.
- AVOID actions that may cause powder to become airborne.
- Use a properly-fitted, NIOSHapproved respirator ("mask"), wear eye protection, and minimize skin contact when responding to a situation where small amounts of suspected fentanyl<sup>†</sup> are visible and may become airborne.
- Follow your department guidelines if the scene involves large amounts of suspected fentanyl<sup>†</sup> (e.g., distribution/storage facility, pill milling operation, clandestine lab, gross contamination, spill or release).

#### When exposure occurs

- Prevent further contamination and notify other first responders and dispatch.
- Do not touch your eyes, mouth, nose or any skin after touching any potentially contaminated surface.
- Wash skin thoroughly with cool water, and soap if available. Do NOT use hand sanitizers as they may enhance absorption.
- Wash your hands thoroughly after the incident and before eating, drinking, smoking, or using the restroom.
- If you suspect your clothing, shoes, and PPE may be contaminated, follow your department guidelines for decontamination.

### If you or other first responders exhibit

- Slow Breathing or No Breathing
- Drowsiness or Unresponsiveness - Constricted or Pinpoint Pupils
- Move away from the source of
- exposure and call EMS.
  Administer naloxone according to your department protocols. Multiple doses may be required.
- If naloxone is not available, rescue breathing can be a lifesaving measure until EMS arrives. Use standard basic life support safety precautions (e.g., pocket mask, gloves) to address the exposure risk.
- If needed, initiate CPR until EMS arrives.



- Collaborative Support From:
  - American College of Medical Toxicologists
     American Industrial Hygiene Association
     Association of State and Territorial Health
  - Officials

    Association of State Criminal Investigative
  - Agencies

    Fraternal Order of Police
- International Association of Chiefs of Police
  International Association of Fire Chiefs
- International Association of Fire Chiefs
   International Association of Fire Fighters
- Major Cities Chiefs Association
- Major County Sheriffs of America
- National Alliance of State Drug Enforcement Agencies
- National Association of Counties
  National Association of County and City
- Health Officials
- National Association of Emergency Medical Technicians
- National Association of EMS Physicians
   National Association of State EMS Officials
- National Governor's Association
  National HIDTA Directors Association
- National HIDTA Directors Association
   National Narcotic Officers' Associations' Coalition
- National Sheriffs' Association
- National Volunteer Fire Council
- Police Executive Research Forum
  Police Foundation
- https://www.whitehouse.gov/ondcp/key-issues/fentanyl

Actions to take ..