

FENTANYL[†]

SAFETY RECOMMENDATIONS FOR FIRST RESPONDERS

[†] For the purposes of this document, fentanyl, related substances, and synthetic opioids (herein after referred to as fentanyl[†]) includes fentanyl analogues (e.g., acetylfentanyl, acrylfentanyl, carfentanil, furanylfentanyl), novel synthetic opioids (e.g., U-47700), and other drugs that may be laced with these substances.

- ▶ **The abuse of drugs containing fentanyl[†] is killing Americans. Misinformation and inconsistent recommendations regarding fentanyl[†] have resulted in confusion in the first responder community.**
- ▶ You as a first responder (law enforcement, fire, rescue, and emergency medical services (EMS) personnel) are increasingly likely to encounter fentanyl[†] in your daily activities (e.g., responding to overdose calls, conducting traffic stops, arrests, and searches).
- ▶ This document provides scientific, evidence-based recommendations to protect yourself from exposure.

WHAT YOU NEED TO KNOW

- ▶ Fentanyl[†] can be present in a variety of forms (e.g., powder, tablets, capsules, solutions, and rocks).
- ▶ Inhalation of airborne powder is MOST LIKELY to lead to harmful effects, but is less likely to occur than skin contact.
- ▶ Incidental skin contact may occur during daily activities but is not expected to lead to harmful effects if the contaminated skin is promptly washed off with water.
- ▶ Personal Protective Equipment (PPE) is effective in protecting you from exposure.
- ▶ Slow breathing or no breathing, drowsiness or unresponsiveness, and constricted or pinpoint pupils are the specific signs consistent with fentanyl[†] intoxication.
- ▶ Naloxone is an effective medication that rapidly reverses the effects of fentanyl[†].

Actions to take . . .

To protect yourself from exposure

- ▶ Wear **gloves** when the presence of fentanyl[†] is suspected.
- ▶ **AVOID actions that may cause powder to become airborne.**
- ▶ Use a properly-fitted, NIOSH-approved **respirator ("mask")**, wear **eye protection**, and minimize skin contact when responding to a situation where small amounts of suspected fentanyl[†] are visible and may become airborne.
- ▶ Follow your department guidelines if the scene involves large amounts of suspected fentanyl[†] (e.g., distribution/storage facility, pill milling operation, clandestine lab, gross contamination, spill or release).

When exposure occurs

- ▶ Prevent further contamination and notify other first responders and dispatch.
- ▶ Do not touch your eyes, mouth, nose or any skin after touching any potentially contaminated surface.
- ▶ Wash skin thoroughly with cool water, and soap if available. **Do NOT use hand sanitizers as they may enhance absorption.**
- ▶ Wash your hands thoroughly after the incident and before eating, drinking, smoking, or using the restroom.
- ▶ If you suspect your clothing, shoes, and PPE may be contaminated, follow your department guidelines for decontamination.

If you or other first responders exhibit

- **Slow Breathing or No Breathing**
- **Drowsiness or Unresponsiveness**
- **Constricted or Pinpoint Pupils**
- ▶ Move away from the source of exposure and call EMS.
- ▶ Administer naloxone according to your department protocols. Multiple doses may be required.
- ▶ If naloxone is not available, rescue breathing can be a lifesaving measure until EMS arrives. Use standard basic life support safety precautions (e.g., pocket mask, gloves) to address the exposure risk.
- ▶ If needed, initiate CPR until EMS arrives.



Collaborative Support From:

- American College of Emergency Physicians
- American College of Medical Toxicologists
- American Industrial Hygiene Association
- Association of State and Territorial Health Officials
- Association of State Criminal Investigative Agencies
- Fraternal Order of Police
- International Association of Chiefs of Police
- International Association of Fire Chiefs
- International Association of Fire Fighters
- Major Cities Chiefs Association
- Major County Sheriffs of America
- National Alliance of State Drug Enforcement Agencies
- National Association of Counties
- National Association of County and City Health Officials
- National Association of Emergency Medical Technicians
- National Association of EMS Physicians
- National Association of State EMS Officials
- National Governor's Association
- National HIDTA Directors Association
- National Narcotic Officers' Associations' Coalition
- National Sheriffs' Association
- National Volunteer Fire Council
- Police Executive Research Forum
- Police Foundation

<https://www.whitehouse.gov/ondcp/key-issues/fentanyl>